Video Article

Measurement of Specific Mycobacterial Mistranslation Rates with Gain-of-function Reporter Systems

Yu-Xiang Chen¹, Miamiao Pan¹, Yue-Meng Chen¹, Babak Javid¹

¹Centre for Global Health and Infectious Diseases, Collaborative Innovation Centre for the Diagnosis and Treatment of Infectious Diseases, Tsinghua University School of Medicine

Correspondence to: Babak Javid at bjavid@gmail.com

URL: <https://www.jove.com/video/59453>

DOI: [doi:10.3791/59453](https://doi.org/10.3791/59453)

Keywords: Immunology and Infection, Issue 146, Mycobacterium, mistranslation, translation fidelity, luciferase, gain-of-function, smegmatis

Date Published: 4/26/2019

Citation: Chen, Y.X., Pan, M., Chen, Y.M., Javid, B. Measurement of Specific Mycobacterial Mistranslation Rates with Gain-of-function Reporter Systems. *J. Vis. Exp.* (146), e59453, doi:10.3791/59453 (2019).

Abstract

The translation of genes into proteins is prone to errors. Although the average rate of translational error in model systems is estimated to be 1/10,000 per codon, the actual error rates vary widely, depending on the species, environment, and codons being studied. We have previously shown that mycobacteria use a two-step pathway for the generation of aminoacylated glutamine and asparagine tRNAs and that this is specifically associated with relatively high error rates due to the modulation of mistranslation rates by an essential component of the pathway, the amidotransferase GatCAB. We modified a previously employed *Renilla*-Firefly dual-luciferase system that had been used to measure mistranslation rates in *Escherichia coli* for use in mycobacteria to measure specific mistranslation rates of glutamate at glutamine codons and aspartate for asparagine codons. Although this reporter system was suitable for the accurate estimation of specific error rates, lack of sensitivity and requirements for excessive manipulation steps made it unsuitable for high-throughput applications. Therefore, we developed a second gain-of-function reporter system, using Nluc luciferase and green fluorescent protein (GFP), which is more amenable to medium/high-throughput settings. We used this system to identify kasugamycin as a small molecule that can decrease mycobacterial mistranslation. Although the reporters that we describe here have been used to measure specific types of mycobacterial mistranslation, they may be modified to measure other types of mistranslation in a number of model systems.

Video Link

The video component of this article can be found at <https://www.jove.com/video/59453/>

Introduction

The flow of information in molecular biology requires the translation of genetic information to functional proteins. As with all biological systems, gene translation also involves measurable errors. Estimates of the rates of error in translation are typically quoted as approximately 1/10,000 per codon (reviewed by Ribas de Pouplana et al.¹). However, error rates vary widely, from fewer than 10⁻⁵ to more than 0.05/codon^{1,2,3,4,5}. The wide range of error rates, spanning more than three orders of magnitude, is due to the fact that errors can arise from multiple steps in the translation pathway: from stochastic, mutational, or stress-induced errors in aminoacylation^{6,7,8,9,10}, physiological misacylation of asparaginyl- and glutaminyl-tRNAs⁵, or ribosomal decoding errors^{2,3,11}. Measurably high error rates, representing over 0.01/codon, suggested that translational errors may perform physiological functions^{1,12} and that mistranslation may be context-specific¹³.

We and others have shown that naturally occurring errors in gene translation may be adaptive, especially during environmental stress^{1,5,12,14,15,16,17,18}. In mycobacteria, errors generated by the two-step indirect glutamine/asparagine tRNA aminoacylation pathway^{19,20} result in a remarkably increased tolerance for the first-line antituberculosis antibiotic rifampicin⁵. Therefore, we speculated that decreasing mycobacterial mistranslation with a small molecule may potentiate killing by rifampicin. We screened for and identified the naturally occurring aminoglycoside kasugamycin as a compound that could decrease mycobacterial mistranslation, potentiate rifampicin-mediated killing of mycobacteria both in vitro and in vivo²¹, and limit the emergence of rifampicin resistance²¹, which threatens the global control of tuberculosis²² — the world's most deadly pathogen.

To study translational error, methods for the measurement of mistranslation have to be employed. There are multiple methods that have been developed for the measurement of mistranslation, each with advantages and disadvantages. Briefly, precision mass spectrometry-based methods have several advantages, the most important of which is that with newer algorithms for the detection of multiple types of translational error, a relatively unbiased measurement of mistranslation can be performed¹⁸. However, mass spectrometry is not very suitable for the measurement of deamidation mistranslation events — precisely the type of mistranslation that occurs in mycobacteria due to the error-prone indirect tRNA aminoacylation pathway. This is because of high-frequency nonenzymatic deamidation that occurs in the processing of samples for mass spectrometry²³, resulting in an extremely high background signal. Therefore, for the detection of errors in this pathway, genetic gain-of-function reporters offer distinct advantages. Specifically, suitable gain-of-function reporters can have extremely low background rates, allowing the measurement of very low error rates¹¹.

Since performing experiments with pathogenic *Mycobacterium tuberculosis* requires specialized facilities and extra precautions, we perform most experiments in the nonpathogenic mycobacterium *M. smegmatis* — and have shown earlier that results between the two species are broadly comparable^{5,21}. To measure mistranslation rates in mycobacteria generated by the indirect tRNA aminoacylation pathway, we modified a *Renilla*-Firefly dual-luciferase system that had been previously developed to measure ribosomal decoding errors in *E. coli*¹¹ for use in mycobacteria. We made three specific modifications: the original reporter did not express efficiently in mycobacteria, and therefore, the sequence was codon-optimized, and the C-terminal three amino acids of the firefly luciferase, serine-lysine-leucine, which has been annotated as a trafficking signal in some systems²⁴, was modified to isoleucine-alanine-valine²⁵. The original reporter had a critical lysine residue in the firefly luciferase mutated. Instead, we mutated either a critically conserved aspartate (D120) or glutamate (E144) residue in the *Renilla* luciferase to asparagine and glutamine, respectively²⁵ (**Figure 1**). The reporter was subcloned into the episomal tetracycline-inducible plasmid pUV-tetOR (see the **Table of Materials**). Gain-of-function reporters mutate critically conserved functional residues in enzymes/fluorescent proteins that renders them nonfunctional^{11,26}. Translational (or theoretically, transcriptional) errors that synthesize the functional variant of the protein would result in measurable enzyme activity in a subset of translated proteins. To correct for variation in protein abundance, the mutated reporter is coexpressed with a functional protein that acts as a benchmark and allows for accurate quantification of gain-of-function¹¹. Whilst the *Renilla*-Firefly dual-luciferase reporter allowed for the accurate measurement of specific mycobacterial mistranslation rates^{5,25} (**Figure 2** and section 1 of the protocol), we quickly realized that it is not suitable for medium/high-throughput screening of molecules that would alter the mistranslation rate. This is due mainly to two reasons, namely, a) the relative lack of potency of *Renilla* luciferase meant that a minimum of 1 mL of mycobacterial culture/sample was required to measure mistranslation rates, and b) the requirement for lysis of the cells prior to enzyme activity measurement required excessive manual handling: mycobacterial cells have a thick and multi-layered cell wall and envelope that is relatively resistant to lysis. Therefore, we looked to develop a new gain-of-function reporter system that could be used with small volumes (e.g., in a 96-well plate system) and did not require cell-lysis for measurements. We used the highly potent Nluc luciferase and identified a critical aspartate residue that, when mutated to asparagine, resulted in 2 logs loss of function (**Figure 1**). Furthermore, the small size of Nluc allowed it to have an N-terminal secretion signal tag — from antigen 85A, a major secreted antigen in mycobacteria²⁷ — that would allow Nluc to be secreted into culture supernatant and circumvent the requirement for cell lysis. The benchmark protein, GFP, was expressed from the same promoter as the mutated Nluc, but from an integrated vector (see the **Table of Materials**), and could be measured in intact cells²¹ (**Figure 3**). Despite these advantages, the Nluc/GFP reporter (section 2 of the protocol) also has disadvantages: the relatively modest reduction in Nluc activity (100-fold) with the D140N mutation would not permit the measurement of extremely low mistranslation rates, making the reporter more suitable as a screening tool than for the accurate measurements of translational error. Furthermore, Nluc has no critical glutamate residues; therefore, only asparagine-to-aspartate error rates could be measured. The general principles described in this work should allow researchers to either use these reporters as we have done or modify reporters as appropriate for an accurate and/or facile measurement of other specific translational error rates in their model system of choice.

Protocol

1. *Renilla*-Firefly Dual-luciferase Reporter

NOTE: For a visual representation of this method, see **Figure 2**.

1. Inoculate mycobacterial reporter strains from -80 °C stock with 2 mL of 7H9 medium. Wild-type dual-luciferase, as well as mutated *Renilla* (reporter) strains, needs to be used to allow the calculation of mistranslation rates (see step 1.7). Shake at 37 °C for 1 to 2 days till OD₆₀₀ reaches the stationary phase (OD > 3).
2. Aliquot and dilute to OD_{600nm} around 0.1 - 0.5. For typical experiments, use three independent biological replicate cultures. Anhydrotetracycline (ATC), a tetracycline analog inducer of reporter expression (which is controlled by a tetracycline-inducible promoter) to a final concentration of 50 ng/mL.
3. To measure effects of kasugamycin on mistranslation rates²¹, add different doses of kasugamycin to the culture (see **Figure 4** for the indicated doses) at the same time (at the tested doses, kasugamycin has no antimicrobial activity). Note that it is important to include at least one non-induced control for each reporter. Culture-induce cultures for 4-6 h at 37 °C with shaking.
4. Transfer the bacterial cultures to a 2 mL tube, and centrifuge at 3,220 x g for 5 min at room temperature to pellet down the bacteria. Discard the supernatant.
5. Disrupt the bacteria by adding 40 µL of 1x passive lysis buffer (supplied by a dual-luciferase kit), which has been diluted (1:1) in double-distilled water. Transfer the resuspended bacterial lysate to a white 96-well plate, one well per sample, and shake at room temperature for 20 min.
CAUTION: Do not over incubate the bacteria in lysis buffer (for more than 30 min).
6. Add 80 µL of firefly substrate to each well, shake for 15 s, and measure the luminescence by luminometer with 1,000 ms as integration time. Use either an automated injector or a multichannel pipette to avoid pipetting errors.
7. Add 80 µL of *Renilla* substrate to each well, shake for 15 s, and measure the luminescence by luminometer with 1,000 ms as integration time.
8. Subtract the background luminescence - measured using either wild-type *M. smegmatis* (i.e., not containing reporters) or uninduced reporter lysate - from the measured values. Use the corrected values to calculate the mistranslation rates of each condition by using the following equation^{11,25}, where DN refers to the activity in the mutated reporter strain:

$$\text{Mistranslation rate} = \frac{\left(\frac{\text{Renilla}}{\text{Firefly}}\right)^{\text{DN}}}{\left(\frac{\text{Renilla}}{\text{Firefly}}\right)^{\text{WT}}}$$

2. Nluc/GFP Reporter

NOTE: For a visual representation of this method, see **Figure 3**.

1. Inoculate the bacterial reporter strain from -80 °C stock with 2 mL of 7H9 medium. Shake at 37 °C for 1 to 2 days till OD₆₀₀ reaches the stationary phase (OD >3).
2. Subculture to 50 mL of 7H9 medium and grow till OD₆₀₀ reaches the late stationary phase (>4).
3. Before aliquoting the bacteria to a 96-well plate, add ATC at a final concentration of 50 ng/mL and mix well. Induction of the bulk culture ensures that all wells contain the same amount of inducer and that the induction of the reporter is synchronized. Aliquot the bacteria to a clear, round-bottomed 96-well plate, with 100 µL of volume in each well.
4. To screen for small molecules affecting mistranslation rates, add the compound at the indicated concentration to select wells. For the purposes of this protocol, use kasugamycin (at doses indicated in **Figure 5**) as an illustration. Add different doses of kasugamycin to select wells (each experimental group should contain at least two biological replicates).
5. Shake and induce the samples at 37 °C for 16-20 h.
NOTE: It is necessary to seal the plate with film. Also, all the edge wells need to be filled with at least 200 µL of sterile water to limit evaporation from the test wells.
6. Take 80 µL from each well, using a multichannel pipette, and transfer the samples to a black 96-well plate (which maximizes the fluorescence signal measurement). Measure the GFP signal by luminometer with 20 ms as integration time.
7. After measuring the GFP signal, centrifuge the plate at 3,220 x g for 10 min. Transfer 50 µL of the supernatant to a white-bottomed 96-well plate (which maximizes the luminescence signal measurement), add 50 µL of Nluc substrate to each well, mix them well, and measure the luminescence by luminometer with 1,000 ms as integration time.
8. Determine the Nluc/GFP ratio by dividing the corrected Nluc luminescence values by GFP fluorescence: this measure (in arbitrary units) is a relative measure of aspartate for asparagine mistranslation.

Representative Results

A cartoon illustrating the general outline of the two reporter systems used in this work are shown in **Figure 1**. An overview of section 1 of the protocol is shown in **Figure 2** and an overview of section 2 in **Figure 3**. The effect of kasugamycin on mycobacterial mistranslation is shown in **Figure 4**, as measured by the *Renilla*-Firefly reporter system. To show the specificity of the kasugamycin action, the reporter was also expressed in a strain of *M. smegmatis* in which *ksgA* was deleted ($\Delta ksgA$). KsgA is an rRNA dimethyl transferase, and *ksgA*-deleted strains are relatively resistant to kasugamycin. Kasugamycin action on mistranslation was also measured using the Nluc/GFP reporter, as shown in **Figure 5**.

Figure 1: Cartoon illustrating the two gain-of-function reporter systems. (A) The *Renilla*-Firefly reporter system is comprised of two luciferase enzymes expressed as a fusion protein and under the control of a tetracycline-inducible promoter. Expression of the wild-type dual enzyme results in high measurable activity of both *Renilla* and firefly luciferases (top). Mutation of a critical aspartate residue, D120, in *Renilla* to asparagine renders the *Renilla* enzyme (D120N) inactive, but firefly luciferase is still active. Mistranslation of asparagine to aspartate (in this case, from physiologically misacylated Asp-tRNA^{Asn} tRNA^{5,21}) results in a small proportion of the translated *Renilla* proteins gaining activity, which can be measured. Comparison of the *Renilla*/Firefly activity of the mutated reporter compared with the wild-type dual enzyme allows the calculation of the asparagine to an aspartate mistranslation rate. (B) The Nluc/GFP reporter system operates on a similar principle. The mutated Nluc (D140N) gene has an N-terminal secretion signal derived from antigen 85A, a major secreted mycobacterial protein. Both the *nluc* and *gfp* genes are expressed from identical tetracycline-inducible promoters, to allow the use of GFP as a relative expression benchmark. Note the lack of wild-type Nluc control strain: this reporter is primarily used in screening, where measurements of relative mistranslation rates are sufficient for the primary screen. [Please click here to view a larger version of this figure.](#)

Figure 2: Outline of measuring the mistranslation rate using *Renilla*-*Firefly* reporter (section 1 of the protocol). Fresh cultures of *M. smegmatis*, transformed with a plasmid expressing the dual-luciferase reporters, are grown. The reporter expression is induced, and investigational compounds or conditions are tested. Following 4–6 h of reporter expression, the cultures are pelleted and lysed and the lysates are tested for dual-luciferase activity. [Please click here to view a larger version of this figure.](#)

Figure 3: Outline of measuring the mistranslation rate Nluc-GFP reporter (section 2 of the protocol). A fresh culture of *M. smegmatis*, transformed with the Nluc and GFP reporters, is grown to a sufficient volume for all the plates to be tested (assuming ~10 mL/96-well plate). Immediately prior to pipetting the bacterial culture into each well, induce the expression of the reporters with ATC added to the bulk culture. Incubate the wells by shaking them overnight. To measure the relative mistranslation rates, transfer the cultures to a black plate (to increase the sensitivity of fluorescence detection), measure the GFP fluorescence, and then, spin the plates down to pellet the bacteria. Transfer the supernatant to a white plate for Nluc activity measurement. [Please click here to view a larger version of this figure.](#)

Figure 4: Representative result of using *Renilla*-Firefly reporter measuring the mistranslation rate in the presence of kasugamycin in different strains. Mistranslation rates of aspartate for asparagine in (A) wild-type *M. smegmatis* and (B) a strain deleted for *ksgA* ($\Delta ksgA$) following treatment with kasugamycin (ksg) as measured by the *Renilla*-Firefly dual reporter. The cultures were treated with kasugamycin at indicated doses (in micrograms/milliliter) for 6 h prior to cell lysis and measurements. The corrected Ren/FF ratios (y-axes) are indicative of the aspartate for asparagine mistranslation rates. The deletion of *ksgA* results in a higher baseline mistranslation rate, which is more resistant to modulation by kasugamycin. The bars indicate the mean values, with standard deviation as error. [Please click here to view a larger version of this figure.](#)

Figure 5: Representative result of using *Nluc*-GFP reporter measuring the mistranslation rate in the presence of kasugamycin. Relative mistranslation rates of aspartate for asparagine in wild-type *M. smegmatis* as measured by the *Nluc*/GFP reporter, following treatment with kasugamycin (ksg) overnight (16 h). The bars indicate the mean values, with standard deviation as error. [Please click here to view a larger version of this figure.](#)

Discussion

The protocol described here can be adapted for the measurement of mistranslation rates in a wide variety of organisms. There are a number of considerations that should be kept in mind when adapting the protocol to other systems. First, the purpose of the measurement should be considered. For the accurate measurement of mistranslation rates using gain-of-function reporters, what is needed is (a) a robust readout assay (e.g., enzymatic function [in this case, luciferase activity] with a broad linear range). Also, look for (b) a loss of function mutation in a critical residue of the reporter protein that results in a very significant loss of function, below the expected rate of mistranslation. For example, if the expected mistranslation rate to be measured is approximately 10^{-3} /codon, the loss of function mutation needs to be greater than 1,000-fold; otherwise, the lower range of mistranslation events will not be sensitively detected. Lastly, for the accurate measurement of mistranslation rates using gain-of-function reporters, (c) a robust benchmark reporter is needed — in this case, firefly luciferase for section 1 of the protocol. Ideally, the benchmark reporter should be fused to the mutated enzymatic reporter, guaranteeing (for all intents and purposes) that they are both expressed in equimolar ratios. The benchmark (and primary reporter) readout should be robust to perturbations to exposed environments. For example, the fluorescence of wild-type GFP is highly susceptible to perturbation by changes in pH²⁸, making it unsuitable as a benchmark for the measurement of mistranslation rates in phagocytosed mycobacteria in macrophages, which reside in phagosomes that are below pH

⁷²⁹. On the other hand, for applications such as small molecule screening, the most important considerations for a primary screen will be the reproducibility of the measurements (i.e., precision, as opposed to accuracy), the minimization of manual handling, and small culture volumes. The accurate measurements of mistranslation rates are less important and can be performed as secondary assays. Finally, it should be noted that the reporters described in this protocol, based on the enzymatic function of luciferase enzymes, will only measure mean mistranslation rates of a bacterial population but cannot give information about single-cell heterogeneity variation in mistranslation rates. Given the importance of single-cell variability in adaptive phenotypes^{30,31,32}, fluorescent reporters for the measurement of mistranslation events have been developed^{12,33}, including for the measurement of mistranslation in mycobacteria⁵.

Having considered the requirements for the measurement of mistranslation, it should be noted that the genetic gain-of-function reporters such as described in this protocol can only measure one type of mistranslation (i.e., one amino acid substitution for one codon) per reporter. Therefore, for the measurement of different mistranslation events, multiple reporters are required. For example, for the measurement of mistranslation by ribosomal decoding errors of near-cognate codons by lysyl-tRNA at one position, at least 16 reporters are required^{2,3,11}. High-precision mass spectrometry and bioinformatics allow for the potential identification of many different types of mistranslation events simultaneously¹⁸; however, these methods also come with caveats. In general, they are less accurate than gain-of-function reporters. Furthermore, as stated earlier, they are less suitable for the detection of certain types of mistranslation, namely those that could be conflated with nonenzymatic deamidation²³. Finally, mass spectrometry is unsuitable as a read-out for medium- or high-throughput screening.

For the adaptation of these reporters and protocols for the measurement of mistranslation in other systems, further considerations include a robust expression of the reporter in the desired model system. We initially attempted to use the dual-luciferase system developed by Farabaugh and colleagues in our mycobacterial system without modifications but had no success, due to a lack of reporter expression. Extensive troubleshooting identified that both codon-optimization and a minor sequence modification of the reporters were required to allow for robust expression²⁵ (and see above). It is conceivable that similar adaptations of the reporters would be required for use in other systems.

Finally, consideration should be given to the type of mistranslation event being measured. For example, if there is a need to measure stop-codon readthrough (nonsense suppression), the stop codon's loss of function mutation needs to be introduced within a noncritical region of the primary reporter protein³⁴. Otherwise, only specific nonsense suppression events that regain the critical residue (and, hence, reporter function) will be measured by the assay, which would potentially lead to a substantial underestimation of the true mistranslation rates. There is increasing evidence that translational error plays a possible adaptive role in a large number of organisms^{1,12,13,35}. However, the measurement of mistranslation events is still limited to a small, albeit growing number of model species. The adaptation of sensitive methods to measure mistranslation has the potential to further increase scientists' understanding of the role of translation error in physiology and pathology.

Disclosures

The authors have nothing to disclose.

Acknowledgments

This work was in part supported by grants from the Bill and Melinda Gates Foundation (OPP1109789), the National Natural Science Foundation of China (31570129), and start-up funds from the Tsinghua University School of Medicine to B.J. B.J. is a Wellcome Trust Investigator (207487/Z/17/Z).

References

1. Ribas de Pouplana, L., Santos, M. A., Zhu, J. H., Farabaugh, P. J., Javid, B. Protein mistranslation: friend or foe? *Trends in Biochemical Sciences*. **39** (8), 355-362 (2014).
2. Leng, T., Pan, M., Xu, X., Javid, B. Translational misreading in *Mycobacterium smegmatis* increases in stationary phase. *Tuberculosis (Edinburgh)*. **95** (6), 678-681 (2015).
3. Manickam, N., Nag, N., Abbasi, A., Patel, K., Farabaugh, P. J. Studies of translational misreading in vivo show that the ribosome very efficiently discriminates against most potential errors. *RNA*. **20** (1), 9-15 (2014).
4. Netzer, N. et al. Innate immune and chemically triggered oxidative stress modifies translational fidelity. *Nature*. **462** (7272), 522-526 (2009).
5. Su, H. W. et al. The essential mycobacterial amidotransferase GatCAB is a modulator of specific translational fidelity. *Nature Microbiology*. **1** (11), 16147 (2016).
6. Li, L. et al. Naturally occurring aminoacyl-tRNA synthetases editing-domain mutations that cause mistranslation in *Mycoplasma* parasites. *Proceedings of the National Academy of Sciences of the United States of America*. **108** (23), 9378-9383 (2011).
7. Li, L. et al. Leucyl-tRNA synthetase editing domain functions as a molecular rheostat to control codon ambiguity in *Mycoplasma* pathogens. *Proceedings of the National Academy of Sciences of the United States of America*. **110** (10), 3817-3822 (2013).
8. Ling, J., Soll, D. Severe oxidative stress induces protein mistranslation through impairment of an aminoacyl-tRNA synthetase editing site. *Proceedings of the National Academy of Sciences of the United States of America*. **107** (9), 4028-4033 (2010).
9. Raina, M. et al. Reduced amino acid specificity of mammalian tyrosyl-tRNA synthetase is associated with elevated mistranslation of Tyr codons. *Journal of Biological Chemistry*. (2014).
10. Wu, J., Fan, Y., Ling, J. Mechanism of oxidant-induced mistranslation by threonyl-tRNA synthetase. *Nucleic Acids Research*. **42** (10), 6523-6531 (2014).
11. Kramer, E. B., Farabaugh, P. J. The frequency of translational misreading errors in *E. coli* is largely determined by tRNA competition. *RNA*. **13** (1), 87-96 (2007).
12. Evans, C. R., Fan, Y., Weiss, K., Ling, J. Errors during Gene Expression: Single-Cell Heterogeneity, Stress Resistance, and Microbe-Host Interactions. *mBio*. **9** (4) (2018).
13. Mohler, K., Ibba, M. Translational fidelity and mistranslation in the cellular response to stress. *Nature Microbiology*. **2**, 17117 (2017).

14. Bullwinkle, T. J. et al. Oxidation of cellular amino acid pools leads to cytotoxic mistranslation of the genetic code. *Elife*. **3** (2014).
15. Fan, Y. et al. Heterogeneity of Stop Codon Readthrough in Single Bacterial Cells and Implications for Population Fitness. *Molecular Cell*. **67** (5), 826-836 e825 (2017).
16. Fan, Y. et al. Protein mistranslation protects bacteria against oxidative stress. *Nucleic Acids Research*. **43** (3), 1740-1748 (2015).
17. Schwartz, M. H., Pan, T. Temperature dependent mistranslation in a hyperthermophile adapts proteins to lower temperatures. *Nucleic Acids Research*. **44** (1), 294-303 (2016).
18. Schwartz, M. H., Waldbauer, J. R., Zhang, L., Pan, T. Global tRNA misacylation induced by anaerobiosis and antibiotic exposure broadly increases stress resistance in *Escherichia coli*. *Nucleic Acids Research*. (2016).
19. Curnow, A. W. et al. Glu-tRNA^{Gln} amidotransferase: a novel heterotrimeric enzyme required for correct decoding of glutamine codons during translation. *Proceedings of the National Academy of Sciences of the United States of America*. **94** (22), 11819-11826 (1997).
20. Rathnayake, U. M., Wood, W. N., Hendrickson, T. L. Indirect tRNA aminoacylation during accurate translation and phenotypic mistranslation. *Current Opinion in Chemical Biology*. **41**, 114-122 (2017).
21. Chaudhuri, S. et al. Kasugamycin potentiates rifampicin and limits emergence of resistance in *Mycobacterium tuberculosis* by specifically decreasing mycobacterial mistranslation. *eLife*. **7** (2018).
22. Toosky, M., Javid, B. Novel diagnostics and therapeutics for drug-resistant tuberculosis. *British Medical Bulletin*. **110** (1), 129-140 (2014).
23. Robinson, N. E., Robinson, A. B. Deamidation of human proteins. *Proceedings of the National Academy of Sciences of the United States of America*. **98** (22), 12409-12413 (2001).
24. Miura, S. et al. Urate oxidase is imported into peroxisomes recognizing the C-terminal SKL motif of proteins. *European Journal of Biochemistry*. **223** (1), 141-146 (1994).
25. Javid, B. et al. Mycobacterial mistranslation is necessary and sufficient for rifampicin phenotypic resistance. *Proceedings of the National Academy of Sciences of the United States of America*. **111** (3), 1132-1137 (2014).
26. Wong, S. Y. et al. Functional role of methylation of G518 of the 16S rRNA 530 loop by GidB in *Mycobacterium tuberculosis*. *Antimicrobial Agents and Chemotherapy*. **57** (12), 6311-6318 (2013).
27. Wiker, H. G., Harboe, M. The antigen 85 complex: a major secretion product of *Mycobacterium tuberculosis*. *Microbiological Reviews*. **56** (4), 648-661 (1992).
28. Roberts, T. M. et al. Identification and Characterisation of a pH-stable GFP. *Scientific Reports*. **6**, 28166 (2016).
29. Li, H., Wu, M., Shi, Y., Javid, B. Over-Expression of the Mycobacterial Trehalose-Phosphate Phosphatase OtsB2 Results in a Defect in Macrophage Phagocytosis Associated with Increased Mycobacterial-Macrophage Adhesion. *Frontiers in Microbiology*. **7**, 1754 (2016).
30. Aldridge, B. B. et al. Asymmetry and aging of mycobacterial cells lead to variable growth and antibiotic susceptibility. *Science*. **335** (6064), 100-104 (2012).
31. Rego, E. H., Audette, R. E., Rubin, E. J. Deletion of a mycobacterial divisome factor collapses single-cell phenotypic heterogeneity. *Nature*. **546** (7656), 153-157 (2017).
32. Zhu, J. H. et al. Rifampicin can induce antibiotic tolerance in mycobacteria via paradoxical changes in *rpoB* transcription. *Nature Communications*. **9** (1), 4218 (2018).
33. Lant, J. T. et al. Visualizing tRNA-dependent mistranslation in human cells. *RNA Biology*. **15** (4-5), 567-575 (2018).
34. Grentzmann, G., Ingram, J. A., Kelly, P. J., Gesteland, R. F., Atkins, J. F. A dual-luciferase reporter system for studying recoding signals. *RNA*. **4** (4), 479-486 (1998).
35. Melnikov, S. V., van den Elzen, A., Stevens, D. L., Thoreen, C. C., Soll, D. Loss of protein synthesis quality control in host-restricted organisms. *Proceedings of the National Academy of Sciences of the United States of America*. (2018).